

The Conserved Kinases CDK-1, GSK-3, KIN-19, and MBK-2 Promote OMA-1 Destruction to Regulate the Oocyte-to-Embryo Transition in *C. elegans*

Masaki Shirayama,^{1,5} Martha C. Soto,^{3,5} Takao Ishidate,^{1,5} Soyoung Kim,^{1,5} Kuniaki Nakamura,¹ Yanxia Bei,¹ Sander van den Heuvel,⁴ and Craig C. Mello^{1,2,*}

¹Program in Molecular Medicine

²Howard Hughes Medical Institute

University of Massachusetts Medical School

373 Plantation Street

Worcester, Massachusetts 01605

³Department of Pathology

UMDNJ-RWJMS

Piscataway, New Jersey 08854

⁴Developmental Biology

Kruytbuilding N305

Padualaan 8

Utrecht University

3584 CH Utrecht

The Netherlands

*Correspondence: craig.mello@umassmed.edu

⁵These authors contributed equally to this work.

(Current Biology 16, 47–55; January 10, 2006)

Figure 6B in the version of this paper originally published online on December 8, 2005 was incorrect because the gradient distribution of MEX-5 in the 1-cell stage was not indicated. The figure has been corrected in the print issue and online, and the authors regret this error.

DOI: 10.1016/j.cub.2005.12.029
